

Lattice Approximation for Stochastic Reaction Diffusion Equations with One-Sided Lipschitz Condition

Martin Sauer^{*†} and Wilhelm Stannat^{*‡}

We consider strong convergence of the finite differences approximation in space for stochastic reaction diffusion equations with multiplicative noise under a one-sided Lipschitz condition only. We derive convergence with an implicit rate depending on the regularity of the exact solution. This can be made explicit if the variational solution has more than its canonical spatial regularity. As an application, spatially extended FitzHugh-Nagumo systems with noise are considered.

1 Introduction

Stochastic partial differential equations arise as a model in many scientific fields and the for applications much needed theory of numerical solutions to such equations is growing successfully. However, it is often the case that particular examples such as stochastic reaction diffusion equations, even with very simple structure of the nonlinear term, fail to fit in the assumptions of the majority of publications in this research area. In this article, we study such an SPDE of reaction diffusion type where the nonlinear drift term satisfies a one-sided Lipschitz condition only and is of the form

$$\left. \begin{aligned} \dot{v}(t, \xi) &= \partial_{\xi}^2 v(t, \xi) + \phi_1(\xi, v(t, \xi), w(t, \xi)) + b(\xi, v(t, \xi), w(t, \xi))\eta(t, \xi) \\ \dot{w}(t, \xi) &= \phi_2(\xi, v(t, \xi), w(t, \xi)). \end{aligned} \right\} \quad (1)$$

for $t \geq 0$, $\xi \in (0, 1)$, where $\eta(t, \xi)$ is some Gaussian white noise. This system consists of two variables v and w , where the former satisfies some semi-linear stochastic evolution equation and the latter is a kind of control variable coupled to v in an equation without diffusion and

^{*}Institut für Mathematik, Technische Universität Berlin, Straße des 17. Juni 136, D-10623 Berlin, Germany

[†]Corresponding author, email: sauer@math.tu-berlin.de

[‡]Bernstein Center for Computational Neuroscience, Philippstr. 13, D-10115 Berlin, Germany

Acknowledgement: This work is supported by the BMBF, FKZ 01GQ1001B.

Date: Berlin, January 23, 2013.

AMS Subject Classification: 60H15, 35R60, 60H35, 65C30, 92C20

Keywords and phrases: Stochastic reaction diffusion equations, Finite difference approximation, FitzHugh-Nagumo system

no further independent external noise. The motivation for such a system comes from the field of neurobiology, in particular we consider the spatially extended stochastic FitzHugh-Nagumo system, modeling the propagation of the action potential in the axon of a neuron, see e. g. [2, 12] and Section 6 for details.

The goal of this article is to prove $\mathcal{L}^p$ -convergence including explicit error estimates of an easy to implement spatial approximation scheme. We consider the well-known finite difference method which has been studied by many authors, but to the best of our knowledge there is no convergence result which gives explicit rates if the nonlinear drift term is not Lipschitz continuous. Crucial references in the literature studying similar equations to (1) are Gyöngy [5], where uniform convergence in probability without a rate is proven, as well as Pettersson & Signahl [15] for a similar result with multiplicative noise. Both results are obtained without a global Lipschitz condition. Moreover we want to mention a series of articles by Gyöngy & Millet [6, 7], where the authors work in the variational approach and incorporate various spatial approximation schemes for SPDEs driven by a finite dimensional noise. However, equation (1) is beyond their results since it excludes polynomially growing nonlinearities. On the other hand, it is worth mentioning that there are existing results by Jentzen [11] and Liu [13] which incorporate our one-sided Lipschitz assumption, but use a different approach, the spectral Galerkin approximation, to show path-wise convergence and $\mathcal{L}^p$ -convergence with explicit rates, respectively.

The SPDE (1) is studied within the variational approach, see e. g. [16], and we only use the a priori known regularity of the exact solution to prove $\mathcal{L}^p$ -convergence with an implicitly given rate in Theorem 3.1. This rate is given in terms of the regularity of the exact solution. If there is a better a priori information on the exact solution, this rate can be made explicit in terms of the approximation parameter. The proof is essentially based on uniform exponential a priori estimates for the approximating solutions obtained in Proposition 4.3 and the Itô formula for the variational solution.

Most of the above mentioned articles concern fully discrete schemes, i. e. a discretization of the space and time domain. We only consider the spatial approximation part, which reduces the infinite dimensional problem to an SDE on a finite dimensional space, and then has to be solved with existing theory and the errors accumulate. Nevertheless, let us make a brief comment on the time discretization. One of the most popular approaches is the Euler-Maruyama scheme because of its simplicity and low computational effort. It is known that this scheme converges to the exact solution of the SDE for equations with globally Lipschitz continuous drift and diffusion coefficients. Of course, also our approximated equation fails to satisfy such a condition. For such equations with super-linearly growing drift and/or diffusion it was shown recently by Hutzenthaler, Jentzen & Kloeden [9] that the Euler-Maruyama approximations diverge in a strong and weak sense. An obvious remedy for this problem is to use e. g. the implicit Euler scheme, which is known to converge, however it requires more computational effort than the explicit scheme. Again Hutzenthaler, Jentzen & Kloeden [10] suggest a so-called tamed Euler scheme that is explicit and computationally less expensive. It is also observed that the runtime as a function of the dimension grows linearly and quadratic for the tamed Euler and implicit Euler scheme, respectively. Of course, this favors the tamed Euler scheme for our high dimensional approximating problem. Thus, we propose solving (1) with a combination of finite differences and the tamed Euler scheme.

The article is structured as follows. In the next section, we describe the precise setting and assumptions on the coefficients of (1) and state the existence and uniqueness result for its corresponding abstract stochastic evolution equation in the sense of variational solutions. In Section 3 we introduce the approximation scheme and state the main result in Theorem 3.1. Sections 4 and 5 contain the proof of this theorem. The last part is devoted to the application of our results to a spatially extended FitzHugh-Nagumo system with noise. This model has been studied by Tuckwell [18] concerning the impact of noise on the generation of action potentials and the reliability of faithful signal transmission. Theorem 3.1 now allows to explicitly quantify the approximation error of the finite difference approximation. As an illustration for the use of our result, we state an estimator for the probability of propagation failure based on numerical observations as well as confidence intervals for this estimators. These depend on the statistical Monte-Carlo and the numerical approximation error and are explicitly constructed.

2 Mathematical Setting and Assumptions

Let $(H, \|\cdot\|_H) = L^2(0, 1)$ with scalar product denoted by $\langle \cdot, \cdot \rangle$ and consider the Laplacian as a linear operator on H , $A : D(A) \subset H \rightarrow H$, i. e.

$$(Av)(\xi) := \partial_\xi^2 v(\xi)$$

equipped with (homogeneous) Neumann boundary conditions in 0 and 1. Corresponding to this, recall the definition of the fractional Sobolev spaces $H^\theta(0, 1)$ as $D((-A)^{\frac{\theta}{2}})$ or for $\theta > 0$ with the equivalent Sobolev-Slobodeckij norm, cf. [17],

$$\|u\|_{H^\theta} := \sum_{s=0}^{\lfloor \theta \rfloor} \|\partial_\xi^s u\|_H + \left(\iint_0^1 \frac{|\partial_\xi^{\lfloor \theta \rfloor} u(\xi) - \partial_\xi^{\lfloor \theta \rfloor} u(\zeta)|^2}{|\xi - \zeta|^{2\theta+1}} d\zeta d\xi \right)^{\frac{1}{2}}.$$

Define in particular the space $(V, \|\cdot\|_V)$ as H^1 with norm $\|u\|_V^2 := \int_0^1 |\partial_\xi u(\xi)|^2 d\xi$ and denote by V^* its topological dual, thus we have the Gelfand triplet $V \hookrightarrow H \hookrightarrow V^*$ with continuous and dense embeddings. As auxiliary spaces we define the product spaces $\mathcal{V} := V \times H$, $\mathcal{H} := H \times H$ with $\mathcal{V} \hookrightarrow \mathcal{H} \hookrightarrow \mathcal{V}^*$ continuously and densely. In the following, we define the abstract drift and diffusion operators mapping $\mathcal{V}$ to $\mathcal{V}^*$.

It is well known that A can be uniquely extended to $A : V \rightarrow V^*$ via an integration by parts

$$_{V^*} \langle Au, v \rangle_V = \int_0^1 \partial_\xi^2 u(\xi) v(\xi) d\xi = - \int_0^1 \partial_\xi u(\xi) \partial_\xi v(\xi) d\xi \leq \|u\| \|v\|.$$

The functions $\phi_1, \phi_2 : [0, 1] \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous and satisfy some one-sided Lipschitz and dissipativity, as well as some (local) Lipschitz and growth condition specified below.

Assumption 1. There exist constants $L, M, \gamma > 0$, $\beta \in \mathbb{R}$ and $1 < m \leq 3$ such that

$$\begin{pmatrix} \phi_1(\xi_1, v_1, w_2) - \phi_1(\xi_2, v_2, w_2) \\ \phi_2(\xi_1, v_1, w_1) - \phi_2(\xi_2, v_2, w_2) \end{pmatrix} \cdot \begin{pmatrix} v_1 - v_2 \\ w_1 - w_2 \end{pmatrix} \leq L(|v_1 - v_2|^2 + |w_1 - w_2|^2), \quad (\text{A1})$$

$$\begin{pmatrix} \phi_1(\xi, v, w) \\ \phi_2(\xi, v, w) \end{pmatrix} \cdot \begin{pmatrix} v \\ w \end{pmatrix} \leq \beta|w|^2 + L|v|^2 - \gamma|v|^{m+1}, \quad (\text{A2})$$

$$|\phi_1(\xi, v, w)| \leq M(1 + |v|^m + |w|), \quad (\text{A3})$$

$$|\phi_2(\xi, v, w)| \leq M(1 + |v| + |w|) \quad (\text{A4})$$

and moreover

$$\begin{aligned} |\phi_1(\xi_1, v_1, w_1) - \phi_1(\xi_2, v_2, w_2)| &\leq L(1 + |v_1|^{m-1} + |v_2|^{m-1}) \\ &\quad \cdot (|\xi_1 - \xi_2| + |v_1 - v_2| + |w_1 - w_2|). \end{aligned} \quad (\text{A5})$$

$$|\phi_2(\xi_1, v_1, w_1) - \phi_2(\xi_2, v_2, w_2)| \leq L(|\xi_1 - \xi_2| + |v_1 - v_2| + |w_1 - w_2|). \quad (\text{A6})$$

for all $\xi, \xi_1, \xi_2 \in [0, 1]$, $v, v_1, v_2, w, w_1, w_2 \in \mathbb{R}$.

A typical form of ϕ_1 is a polynomial in v of odd degree and negative leading coefficient together with a linear perturbation in w , for example $\phi_1(v, w) = v - v^3 - w$ similar as in the FitzHugh-Nagumo system described in Section 6. With these assumptions we can define the Nemytskii operator $\Phi : \mathcal{V} \rightarrow \mathcal{V}^*$ by

$$(\Phi(v, w))(\xi) := \begin{pmatrix} \phi_1(\xi, v(\xi), w(\xi)) \\ \phi_2(\xi, v(\xi), w(\xi)) \end{pmatrix}$$

and estimate for $v_1, v_2 \in V$, $w_1, w_2 \in H$ with the embedding $V \hookrightarrow L^\infty(0, 1)$ as follows

$$\mathcal{V}^* \langle \Phi(v_1, w_1), (v_2, w_2) \rangle_{\mathcal{V}} \leq C \left(\|v_1\|_V^m \|v_2\|_V + (1 + \|w_1\|_H) \|v_2\|_H + (1 + \|w_1\|_H + \|v_1\|_H) \|w_2\|_H \right).$$

Now that the assumptions on the drift of equation (1) are stated, let us focus on the noise process η , formally given as the derivative of some Q -Wiener process $\sqrt{Q}W(t)$, where $(W(t))_{t \geq 0}$ is a cylindrical Wiener process on H w. r. t. an underlying probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, \mathbb{P})$. In order to obtain a solution to (1) (in a sense specified below) we cannot treat the case of space-time white noise, i. e. $Q = I$, but only the case of colored noise with nuclear covariance operator Q . The assumptions on Q and b are combined below.

Assumption 2. Let $\theta \in (\frac{1}{2}, 1)$ and $Q \in L(H)$ be symmetric and positive definite. Assume that $\text{tr}(-A)^\theta Q < \infty$, in particular Q admits an integral kernel of the form

$$(Qv)(\xi) = \int_0^1 q(\xi, \zeta) q(\rho, \zeta) v(\rho) d\zeta d\rho, \quad q \in H^\theta((0, 1)^2). \quad (\text{A7})$$

The scalar valued noise intensity $b : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is supposed to be bounded and Lipschitz continuous, in particular

$$|b(\xi, v)| \leq 1 \quad \text{and} \quad |b(\xi_1, v_1) - b(\xi_2, v_2)| \leq (|\xi_1 - \xi_2| + |v_1 - v_2|) \quad (\text{A8})$$

for all $\xi, \xi_1, \xi_2 \in [0, 1]$ and $v, v_1, v_2 \in \mathbb{R}$.

In the same manner as for the nonlinear drift, we define the Nemytskii operator $(B(v))(\xi) := b(\xi, v(\xi))$. Furthermore, let $x = (v, w)^T \in \mathcal{V}$ and

$$\mathcal{A}(x) := \begin{pmatrix} Av \\ 0 \end{pmatrix} + \Phi(v, w), \quad \mathcal{B}(x) := \begin{pmatrix} B(v)\sqrt{Q} \\ 0 \end{pmatrix}.$$

Then, the stochastic evolution equation for $X(t) := (v(t), w(t))^T$ corresponding to equation (1) is given by

$$dX(t) = \mathcal{A}(X(t)) dt + \mathcal{B}(X(t)) dW(t), \quad X(0) = x_0. \quad (2)$$

Existence and uniqueness of a solution for such an equation can be studied in the variational framework, see for example [16], in particular a recent extension in [14]. The following theorem is a corollary to [14, Theorem 1.1].

Theorem 2.1. *Let $T > 0$, $x_0 \in L^{2m}(\Omega, \mathcal{F}, \mathbb{P}; H)$ and (A1)-(A4), (A7), (A8) are satisfied. Then, equation (2) has a unique variational solution $(X(t))_{t \in [0, T]}$ which satisfies*

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|X(t)\|_{\mathcal{H}}^{2m} + \int_0^T \|X(t)\|_{\mathcal{V}}^2 dt \right] < \infty.$$

Proof. We have to verify the conditions (H1)-(H4) in [14]. Of course, for $x_1, x_2, x_3 \in \mathcal{V}$ the map $s \mapsto \nu^* \langle \mathcal{A}(x_1 + sx_2), x_3 \rangle_{\mathcal{V}}$ is continuous on $\mathbb{R}$, hence (H1) holds. Furthermore, we can easily obtain the monotonicity of $\mathcal{A}$ by the one-sided Lipschitz condition (A1)

$$2\nu^* \langle \mathcal{A}(x_1) - \mathcal{A}(x_2), x_1 - x_2 \rangle_{\mathcal{V}} \leq -2\|v_1 - v_2\| + 2L\|x_1 - x_2\|_{\mathcal{H}}^2.$$

Moreover, $\mathcal{B}$ is Lipschitz continuous in the Hilbert-Schmidt norm since

$$\begin{aligned} \|\mathcal{B}(x_1) - \mathcal{B}(x_2)\|_{\text{HS}} &\leq \|B(v_1) - B(v_2)\|_{L(H^\theta, H)} \|(-A)^{\frac{\theta}{2}} \sqrt{Q}\|_{\text{HS}} \\ &\leq c_\theta \|v_1 - v_2\|_H (\text{tr}(-A)^\theta Q)^{\frac{1}{2}}. \end{aligned} \quad (3)$$

The latter inequality holds because of the embedding $H^\theta(0, 1) \hookrightarrow L^\infty(0, 1)$ for $\theta > \frac{1}{2}$ with constant c_θ . Thus, we have verified (H2) with $\rho \equiv 0$. (A2) together with the fact that $\|B(x)\|_{L(H)} \leq 1$ implies

$$2\nu^* \langle \mathcal{A}(x), x \rangle_{\mathcal{V}} + \|\mathcal{B}(x)\|_{\text{HS}}^2 \leq -2\|x\|_{\mathcal{V}}^2 + 2\max\{L, \beta + 1\}\|x\|_{\mathcal{H}}^2 + \text{tr } Q$$

for all $x \in \mathcal{V}$ and (H2) holds with $\alpha = 2$. With (A3) and (A4) the matching growth condition is

$$\begin{aligned} \|\mathcal{A}(x)\|_{\mathcal{V}^*} &\leq \|x\|_{\mathcal{V}} + \sup_{\|u\|_{\mathcal{V}}=1} \int_0^1 |\phi_1(\xi, v(\xi), w(\xi))u(\xi)| d\xi + \sup_{\|u\|_{H=1}} \int_0^1 |\phi_2(\xi, v(\xi), w(\xi))u(\xi)| d\xi \\ &\leq \|x\|_{\mathcal{V}} + 2M \sup_{\|u\|_{H=1}} \int_0^1 (1 + |w(\xi)| + |v(\xi)|) |u(\xi)| d\xi + M \sup_{\|u\|_{\mathcal{V}}=1} \int_0^1 |v(\xi)|^m |u(\xi)| d\xi \\ &\leq \|x\|_{\mathcal{V}}^2 + 2M(1 + \|w\|_H + \|v\|_H) + c_1 M \|v\|_V \left(\int_0^1 |v(\xi)|^{2m-2} d\xi \right)^{\frac{1}{2}}, \end{aligned}$$

where we used the embedding $V \hookrightarrow L^\infty(0, 1)$. The latter is finite if $m \leq 3$, hence

$$\|\mathcal{A}(x)\|_{\mathcal{V}^*}^2 \leq C(1 + \|x\|_{\mathcal{V}}^2)(1 + \|x\|_{\mathcal{H}}^{2m-2})$$

and condition (H4) in [14] holds. $\square$

3 The Finite Differences Scheme and the Main Result

In this section we briefly describe the approximation scheme and state the main convergence result in Theorem 3.1. Equation (2) is evaluated on a simple grid approximating the domain $(0, 1)$, given by $\frac{1}{n}\{0, \dots, n\}$ and the vectors $v^n, w^n \in \mathbb{R}^{n+1}$ representing the functions v and w on this grid. Furthermore, we introduce the spaces $V_n \cong \mathbb{R}^{n+1}$ with the discrete boundary conditions $u_1 - u_0 = u_n - u_{n-1} = 0$ corresponding to the homogeneous Neumann boundary. We equip V_n with the semi-norm $\|v\|_n^2 := n \sum_{k=1}^n (v_k - v_{k-1})^2$. In order to imitate the variational approach in the discrete setting we want a summation by parts formula to hold, i. e.

$$\frac{1}{n} \sum_{k=0}^n (A^n v)_k u_k = -n \sum_{k=1}^n (v_k - v_{k-1})(u_k - u_{k-1}) \quad (4)$$

for all $v \in V^n$, $u \in \mathbb{R}^{n+1}$. The resulting finite difference approximation A^n of A is just the discrete Laplacian, given by

$$(A^n v)_k := n^2(v_{k+1} - 2v_k + v_{k-1}), \quad 1 \leq k \leq n-1, \quad v \in V_n$$

together with the appropriate boundary values $(A^n v)_0 = n(v_1 - v_0)$, $(A^n v)_n = n(v_n - v_{n-1})$ which are 0 for $v \in V^n$. The nonlinearities are simply evaluated point-wise, i. e.

$$\phi_{1,2}^n(v, w) := \left(\phi_{1,2}\left(\frac{k}{n}, v_k, w_k\right) \right)_{0 \leq k \leq n} \quad \text{and} \quad b^n(v) := \text{diag} \left(b\left(\frac{k}{n}, v_k\right) \right)_{0 \leq k \leq n}$$

for $v, w \in V_n$ where by $\text{diag } v$ we denote the diagonal matrix with the entries of the vector v on the main diagonal. In the next step let us construct the approximating noise in terms of the given realization of the driving cylindrical Wiener process $(W(t))_{t \geq 0}$. Recall that

$$\langle W(t), \sqrt{n} 1_{[\frac{k-1}{n}, \frac{k}{n})} \rangle =: \beta_k(t), \quad 1 \leq k \leq n \quad (5)$$

defines a family of n iid real valued Brownian motions. The spatial covariance structure given by the kernel q is discretized as follows,

$$q_{k,l}^n := n^2 \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} q(\xi, \zeta) d\zeta d\xi, \quad 1 \leq k, l \leq n,$$

together with $q_{0,l}^n := 0$, $1 \leq l \leq n$. With this discrete covariance matrix we can replace q by a piecewise constant kernel, i. e. for $1 \leq k \leq n$ and $\xi \in [\frac{k-1}{n}, \frac{k}{n})$

$$\begin{aligned} \int_0^1 q(\xi, \zeta) W(t, \zeta) d\zeta &\rightsquigarrow \sum_{l=1}^n q_{k,l}^n \int_0^1 1_{[\frac{l-1}{n}, \frac{l}{n})}(\zeta) W(t, \zeta) d\zeta = \frac{1}{\sqrt{n}} \sum_{l=1}^n q_{k,l}^n \langle W(t), \sqrt{n} 1_{[\frac{l-1}{n}, \frac{l}{n})} \rangle \\ &= \frac{1}{\sqrt{n}} \sum_{l=1}^n q_{k,l}^n \beta_l(t) =: (\sqrt{Q^n} P_n W(t))_k, \end{aligned}$$

where $P_n h = (\langle h, \sqrt{n} 1_{[\frac{l-1}{n}, \frac{l}{n})} \rangle)_{1 \leq l \leq n}$. Denote by $W^n(t) = P_n W(t)$ the resulting n -dimensional Brownian motion and the finite dimensional system of stochastic differential equations approximating equation (2) or rather (1) is

$$\left. \begin{aligned} dv^n(t) &= \left[A^n v^n(t) + \phi_1^n(v^n(t), w^n(t)) \right] dt + b^n(v^n(t)) \sqrt{Q^n} dW^n(t), \\ dw^n(t) &= \left[\phi_2^n(v^n(t), w^n(t)) \right] dt. \end{aligned} \right\} \quad (6)$$

Standard results on SDEs imply the existence of a unique strong solution $(v^n(t), w^n(t))$ to (6), see e.g. [16, Chapter 3]. In order to formulate the convergence result, we embed the processes v^n and w^n into the space $C([0, T], V)$ by linear interpolation w.r.t. the space variable ξ . Given $v \in V_n$, define

$$\tilde{v}(\xi) := (n\xi - k + 1)v_k + (k - n\xi)v_{k-1}, \quad \xi \in \left[\frac{k-1}{n}, \frac{k}{n}\right].$$

Denote by $\iota_n : V_n \rightarrow V; v \mapsto \tilde{v}$ the embedding given above. Furthermore define

$$\tilde{X}^n(t) := \begin{pmatrix} \tilde{v}^n(t) \\ \tilde{w}^n(t) \end{pmatrix} \quad \text{and} \quad E^n(t) := X(t) - \tilde{X}^n(t).$$

Theorem 3.1. *Suppose Assumptions 1 and 2 hold, $v_0 \in \mathcal{L}^\infty(\Omega, \mathcal{F}, \mathbb{P}; C([0, 1]))$ and $w_0 \in \mathcal{L}^\infty(\Omega, \mathcal{F}, \mathbb{P}; C^1([0, 1]))$. Then for every $1 \leq p < \infty$ there exists a finite constant $C = C(v_0, w_0, p, T, Q, L, \beta, \gamma)$ such that*

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|E^n(t)\|_{\mathcal{H}}^p \right]^{\frac{1}{p}} \leq C \left(\mathbb{E} \left[\|E^n(0)\|_{\mathcal{H}}^{2p} \right] + \mathbb{E} \left[\left(\int_0^T I_n(v(t))^2 dt \right)^p \right]^{\frac{1}{2}} \right)^{\frac{1}{2p}}$$

which converges to 0 as $n \rightarrow \infty$. The rate of convergence is implicitly given by the expression

$$I_n(v(t)) := \left(\sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left(v'(t, \zeta) - v'(t, \zeta \pm \frac{1}{n}) \right)^2 d\zeta \right)^{\frac{1}{2}}.$$

If in addition $v \in \mathcal{L}^{p^*}(\Omega, \mathcal{F}, \mathbb{P}; L^2([0, T], H^{1+\alpha}))$ for some $\alpha > 0$ and $1 < p^* < \infty$, then the rate of convergence is explicitly given by

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|E^n(t)\|_{\mathcal{H}}^p \right]^{\frac{1}{p}} \leq C \left(\mathbb{E} \left[\|E^n(0)\|_{\mathcal{H}}^{2p} \right]^{\frac{1}{2p}} + n^{-\frac{1}{2} \min\{1, \alpha\}} \right)$$

for all $p < p^*$ and some constant $C = C(v_0, w_0, p, p^*, T, Q, L, \beta, \gamma, \|v\|_{\mathcal{L}^{p^*}(L^2([0, T], H^{1+\alpha}))})$.

4 Uniform A Priori Estimates

In addition to the a priori estimates in Theorem 2.1 the proof of Theorem 3.1 requires such estimates corresponding to the approximating solutions $\tilde{v}^n$ and $\tilde{w}^n$. These estimates are formally obtained by applying Itô's formula to the square of the $\mathcal{H}$ -norm and we will do the same in the discrete case and obtain an exponential a priori estimate for the $L^2([0, T], V)$ -norm of $\tilde{v}^n$ and $L^\infty([0, T], V)$ -norm of $\tilde{w}^n$ uniform in the parameter n . Moreover due to (A2), we have control over the discrete l_{m+1} -norm of v^n uniformly in n . For this purpose define

$$l_p^n(v) := \frac{1}{n} \sum_{k=1}^n |v_k|^p, \quad v \in V_n, \quad 1 \leq p < \infty.$$

Lemma 4.1. *Let $n \in \mathbb{N}$. The approximating solution $(v^n(t), w^n(t))$ satisfies the estimate*

$$\begin{aligned} & \mathbb{E} \left[l_2^n(v^n(t)) + l_2^n(w^n(t)) + 2 \int_0^t \|v^n(s)\|_n^2 ds + 2\gamma \int_0^t l_{m+1}^n(v^n(s)) ds \right] \\ & \leq e^{2 \max\{L, \beta\}T} \left(\mathbb{E} [l_2^n(v^n(0)) + l_2^n(w^n(0))] + T \operatorname{tr} Q \right) \quad \text{for all } t \in [0, T]. \end{aligned} \quad (7)$$

Proof. Itô's formula applied to $(v_k^n)^2 + (w_k^n)^2$, the summation by parts formula (4) and the dissipativity condition on ϕ_1, ϕ_2 in (A2) imply

$$\begin{aligned} & l_2^n(v^n(t)) + l_2^n(w^n(t)) + 2 \int_0^t \|v^n(s)\|_n^2 ds + 2\gamma \int_0^t l_{m+1}^n(v^n(s)) ds \\ & \leq l_2^n(v^n(0)) + l_2^n(w^n(0)) + 2L \int_0^t l_2^n(v^n(s)) ds + 2\beta \int_0^t l_2^n(w^n(s)) ds \\ & \quad + \frac{1}{n^2} \int_0^t \sum_{k,l=1}^n |b(\frac{k}{n}, v_k^n(s)) q_{k,l}^n|^2 ds + \frac{2}{n\sqrt{n}} \int_0^t \sum_{k,l=1}^n b(\frac{k}{n}, v_k^n(s)) v_k^n(s) q_{k,l}^n d\beta_l(s). \end{aligned}$$

With a standard localization argument the stochastic integral vanishes after taking the expectation and since b is bounded we obtain

$$\begin{aligned} & \mathbb{E} \left[l_2^n(v^n(t)) + l_2^n(w^n(t)) + 2 \int_0^t \|v^n(s)\|_n^2 ds + 2\gamma \int_0^t l_{m+1}^n(v^n(s)) ds \right] \\ & \leq \mathbb{E} \left[l_2^n(v^n(0)) + l_2^n(w^n(0)) + 2 \max\{L, \beta\} \int_0^t l_2^n(v^n(s)) + l_2^n(w^n(s)) ds + t \operatorname{tr} Q \right]. \end{aligned}$$

Gronwall's lemma now yields the result. $\square$

Unlike v , the equation for w has no regularizing linear part. Nevertheless, one can improve the a priori estimate significantly if the initial condition has more regularity, because the coupling with v is Lipschitz continuous.

Lemma 4.2. *The following improved a priori estimate holds for every $n \in \mathbb{N}$*

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|w^n(t)\|_n^2 \right] \leq L e^{5 \max\{L, \beta\}T} \left(\mathbb{E} [\|w^n(0)\|_n^2 + l_2^n(v^n(0)) + l_2^n(w^n(0))] + T \operatorname{tr} Q \right). \quad (8)$$

Proof. Consider the equation for $w_k^n(t)$ for itself and a straightforward calculation yields

$$\begin{aligned} & |w_k^n(t) - w_{k-1}^n(t)|^2 = |w_k^n(0) - w_{k-1}^n(0)|^2 \\ & \quad + 2 \int_0^t (w_k^n(s) - w_{k-1}^n(s)) \left(\phi_2\left(\frac{k}{n}, v_k^n(s), w_k^n(s)\right) - \phi_2\left(\frac{k-1}{n}, v_{k-1}^n(s), w_{k-1}^n(s)\right) \right) ds \\ & \leq 2L \int_0^t |w_k^n(s) - w_{k-1}^n(s)|^2 + |w_k^n(s) - w_{k-1}^n(s)| \left(\frac{1}{n} + |v_k^n(s) - v_{k-1}^n(s)| \right) ds \\ & \leq 3L \int_0^t |w_k^n(s) - w_{k-1}^n(s)|^2 ds + 2L \int_0^t \left(\frac{1}{n^2} + |v_k^n(s) - v_{k-1}^n(s)|^2 \right) ds. \end{aligned}$$

Hence, a summation over k and an application of Gronwall's lemma yield

$$\mathbb{E} \left[\sup_{t \in [0, T]} \|w^n(t)\|_n^2 \right] \leq e^{3LT} \left(\|w^n(0)\|_n^2 + 2L \mathbb{E} \left[\int_0^T (1 + \|v^n(t)\|_n^2) dt \right] \right).$$

Now use the estimate (7) to conclude (8). $\square$

By a similar strategy as in Lemmas 4.1 and 4.2 we can also obtain exponential a priori estimates. This is due to a standard decomposition of the exponential of a martingale into a supermartingale and the exponential of its quadratic variation process. The latter can be bounded independent of the size of $\text{tr } Q$ by the estimate in l_{m+1} . The precise statement is the following.

Proposition 4.3. *Let $v_0 \in \mathcal{L}^\infty(\Omega, \mathcal{F}, \mathbb{P}; C([0, 1]))$, $w_0 \in \mathcal{L}^\infty(\Omega, \mathcal{F}, \mathbb{P}; C^1([0, 1]))$ and $\alpha > 0$ arbitrary. Then*

$$\mathbb{E} \left[\exp \left(\alpha \left(\int_0^T (\|\tilde{v}^n(t)\|_V^2 + \gamma l_{m+1}^n(v^n(t))) dt + \sup_{t \in [0, T]} \|\tilde{w}^n(t)\|_V^2 \right) \right) \right] \leq C_{4.3}$$

uniformly in $n \in \mathbb{N}$, where $C_{4.3} = C(\alpha, v_0, w_0, T, Q, L, m)$ is an explicitly known finite constant.

Proof. Note, that the Itô formula in the proof of Lemma 4.1 and Gronwall's lemma imply

$$\int_0^T (\|v^n(t)\|_n^2 + \gamma l_{m+1}^n(v^n(t))) dt \leq \frac{1}{2} e^{2 \max\{L, \beta\}T} \left(l_2^n(v^n(0)) + l_2^n(w^n(0)) + T \text{tr } Q + M_T \right),$$

where M_T denotes the stochastic integral. Also, we have seen in Lemma 4.2 that

$$\sup_{t \in [0, T]} \|w^n(t)\|_n^2 \leq e^{3LT} \left(\|w^n(0)\|_n^2 + 2L \int_0^T \|v^n(t)\|_n^2 dt \right).$$

Hence, a combination of the two inequalities above yields that for any $\alpha > 0$

$$\begin{aligned} & \exp \left(\alpha \left(\int_0^T (\|v^n(t)\|_n^2 + \gamma l_{m+1}^n(v^n(t))) dt + \sup_{t \in [0, T]} \|w^n(t)\|_n^2 \right) \right) \\ & \leq \exp \left(\alpha C_1 (l_2^n(v^n(0)) + l_2^n(w^n(0)) + \|w^n(0)\|_n^2 + T \text{tr } Q) \right) \cdot \exp \left(\alpha C_2 M_T \right). \end{aligned} \tag{9}$$

Here, $C_1 = (2L + 1)e^{5 \max\{L, \beta\}T}$ and $C_2 = e^{2 \max\{L, \beta\}T} + 2Le^{3LT + 2 \max\{L, \beta\}T}$. Now recall that for every continuous local martingale vanishing at $t = 0$ the process

$$Z_t^\alpha := \exp \left(\alpha M_t - \frac{\alpha^2}{2} \langle M \rangle_t \right)$$

is again a continuous local martingale, thus a supermartingale by Fatou's lemma and therefore $\mathbb{E}[Z_t^\alpha] \leq 1$ for all t . With this information we can derive

$$\mathbb{E} \left[\exp \left(\frac{\alpha}{2} M_t \right) \right] \leq \mathbb{E} \left[\exp \left(\frac{\alpha^2}{2} \langle M \rangle_t \right) \right]^{\frac{1}{2}}. \tag{10}$$

Now we can take the expectation in (9), Hölder's inequality and then (10) implies

$$\begin{aligned}
& \mathbb{E} \left[\exp \left(\alpha \left(\int_0^T (\|v^n(t)\|_n^2 + \gamma l_{m+1}^n(v^n(t))) dt + \sup_{t \in [0, T]} \|w^n(t)\|_n^2 \right) \right) \right]^2 \\
& \leq \mathbb{E} \left[\exp \left(2\alpha C_1 (l_2^n(v^n(0)) + l_2^n(w^n(0)) + \|w^n(0)\|_n^2 + T \operatorname{tr} Q) \right) \right] \mathbb{E} \left[\exp \left(2\alpha C_2 M_T \right) \right] \\
& \leq \mathbb{E} \left[\exp \left(2\alpha C_1 (l_2^n(v^n(0)) + l_2^n(w^n(0)) + \|w^n(0)\|_n^2 + T \operatorname{tr} Q) \right) \right] \mathbb{E} \left[\exp \left(8\alpha^2 C_2^2 \langle M \rangle_T \right) \right]^{\frac{1}{2}}.
\end{aligned}$$

The quadratic variation $\langle M \rangle_T$ can be bounded by the integrals on the left hand side, in particular by the l_{m+1}^n -norm of v^n .

$$\begin{aligned}
\langle M \rangle_T &= \frac{4}{n^3} \int_0^T \sum_{k,l=1}^n \left(b\left(\frac{k}{n}, v_k^n(t)\right) v_k^n(t) q_{k,l}^n \right)^2 dt \\
&\leq \frac{4}{n^3} \int_0^T \sum_{k,l=1}^n (v_k^n(t) q_{k,l}^n)^2 dt \\
&\leq \frac{4}{n^3} \int_0^T \left(\sum_{k=1}^n (v_k^n(t))^2 \right) \left(\sum_{k,l=1}^n (q_{k,l}^n)^2 \right) dt \\
&\leq 4 \operatorname{tr} Q \int_0^T l_2^n(v^n(t)) dt \\
&\leq 4 \operatorname{tr} Q \int_0^T \delta l_{m+1}^n(v^n(t)) + \delta^{\frac{m-1}{2}} \left(\frac{2}{\delta(m+1)} \right)^{\frac{m+1}{m-1}} dt,
\end{aligned}$$

where we used Young's inequality in the last line for arbitrary $\delta > 0$. In particular, we can choose $\delta = \gamma(32\alpha C_2^2 \operatorname{tr} Q)^{-1}$ such that

$$\begin{aligned}
& \mathbb{E} \left[\exp \left(\alpha \left(\int_0^T (\|v^n(t)\|_n^2 + \gamma l_{m+1}^n(v^n(t))) dt + \sup_{t \in [0, T]} \|w^n(t)\|_n^2 \right) \right) \right] \\
& \leq e^{\frac{8}{3} C_\delta T \operatorname{tr} Q} \mathbb{E} \left[\exp \left(2\alpha C_1 (l_2^n(v^n(0)) + l_2^n(w^n(0)) + \|w^n(0)\|_n^2 + T \operatorname{tr} Q) \right) \right]^{\frac{2}{3}}, \tag{11}
\end{aligned}$$

where C_δ is the constant appearing in Young's inequality above. Finally, observe that for $v \in V^n$ the piecewise linear interpolation $\tilde{v}$ is in V and its weak derivative is explicitly given by the differences $n(v_k - v_{k-1})$. Hence

$$\|\tilde{v}\|_V^2 = \int_0^1 |\partial_\xi \tilde{v}(\xi)|^2 d\xi = \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} n^2 (v_k - v_{k-1})^2 d\xi = n \sum_{k=1}^n (v_k - v_{k-1})^2.$$

Moreover, note that for $v_0 \in C([0, 1])$ and $w_0 \in C^1([0, 1])$ we can define the point-wise evaluation $v^n(0)$ and $w^n(0)$, which satisfy

$$l_2^n(v^n(0)) \leq \|v_0\|_{C([0,1])}^2 \quad \text{and} \quad l_2^n(w^n(0)) + \|\tilde{w}_0^n(0)\|_V^2 \leq \|w_0\|_{C^1([0,1])}^2$$

and we found an upper bound for (11) uniformly in $n \in \mathbb{N}$. $\square$

5 Error Estimation, Proof of Theorem 3.1

This section is devoted to the proof of Theorem 3.1 and the estimation of the error in $\mathcal{L}^p(\Omega, \mathcal{F}, \mathbb{P}; C([0, T], \mathcal{H}))$. We proceed in several steps, first some path-wise estimates and then using the Burkholder-Davis-Gundy inequality together with the a priori information on the approximating solution $\tilde{X}^n$ in Proposition 4.3 to prove that $\tilde{X}^n$ converges strongly to the exact solution X .

Step 1: Itô's formula applied to the square of the $\mathcal{H}$ -norm (which is available for the variational solution) implies

$$\begin{aligned} \|E^n(t)\|_{\mathcal{H}}^2 - \|E^n(0)\|_{\mathcal{H}}^2 &= 2 \int_0^t \nu^* \langle Av(s) - \iota_n A^n v^n(s), v(s) - \tilde{v}^n(s) \rangle_V ds \\ &\quad + 2 \int_0^t \nu^* \langle \Phi(v(s), w(s)) - \iota_n \Phi^n(v^n(s), w^n(s)), E^n(s) \rangle_V ds \\ &\quad + 2 \int_0^t \langle v(s) - \tilde{v}^n(s), (B(v(s))\sqrt{Q} - \iota_n b^n(v^n(s))\sqrt{Q^n}P_n) dW(s) \rangle \\ &\quad + \int_0^t \|B(v(s))\sqrt{Q} - \iota_n b^n(v^n(s))\sqrt{Q^n}P_n\|_{\text{HS}}^2 ds \\ &= E_1 + E_2 + E_3 + E_4. \end{aligned} \tag{12}$$

Each of the error terms E_1, E_2, E_4 can be split into two parts with the first one only involving the difference E^n and the second one the approximation of the parameters of the equation. For the stochastic integral in E_3 we use the same argument after applying the Burkholder-Davis-Gundy inequality later. Now let us begin with, for instance E_1 where an integration by parts yields

$$E_1 = -2 \int_0^t \|v(s) - \tilde{v}^n(s)\|_V^2 ds + 2 \int_0^t \nu^* \langle A\tilde{v}^n(s) - \iota_n A^n v^n(s), v(s) - \tilde{v}^n(s) \rangle_V ds.$$

Moreover the one-sided Lipschitz condition (A1) for the nonlinear drift part and the Lipschitz continuity of B in (3) yield

$$\begin{aligned} E_2 &\leq 2L \int_0^t \|E^n(s)\|_{\mathcal{H}}^2 ds + 2 \int_0^t \nu^* \langle \Phi(\tilde{v}^n(s), \tilde{w}^n(s)) - \iota_n \Phi^n(v^n(s), w^n(s)), E^n(s) \rangle_V ds \\ E_3 &\leq 2c_\theta \text{tr}(-A)^\theta Q \int_0^t \|E^n(s)\|_{\mathcal{H}}^2 ds + 2 \int_0^t \|B(\tilde{v}^n(s))\sqrt{Q} - \iota_n b^n(v^n(s))\sqrt{Q^n}P_n\|_{\text{HS}}^2 ds. \end{aligned}$$

In the following, we estimate each of the error terms for fixed s , thus we drop the time dependence in the notation.

Step 2: Approximation error of the Laplacian

At first, we will prove that the error term coming from the linear part converges to 0 if $v \in V$, i.e. we need only the guaranteed regularity of the variational solution. The rate of convergence is not uniform and given implicitly in terms of the solution v . Secondly, if we assume additional regularity on v , we deduce an explicit rate in terms of n . Let $\varphi \in V$, then

$$\nu^* \langle A\tilde{v}^n - \iota_n A^n v^n, \varphi \rangle_V = \nu^* \langle A\tilde{v}^n, \varphi \rangle_V - \langle \iota_n A^n v^n, \varphi \rangle$$

since the linear interpolation is in V . In the first term an integration by parts yields

$$V^* \langle A\tilde{v}^n, \varphi \rangle_V = -n \sum_{k=1}^n (v_k^n - v_{k-1}^n) (\varphi(\frac{k}{n}) - \varphi(\frac{k-1}{n})). \quad (13)$$

In the second term we use the summation by parts formula (4), hence

$$\begin{aligned} \langle \iota_n A^n v^n, \varphi \rangle &= \sum_{k=1}^n (A^n v^n)_k \int_{\frac{k-1}{n}}^{\frac{k}{n}} (n\xi - k + 1) \varphi(\xi) d\xi + (A^n v^n)_{k-1} \int_{\frac{k-1}{n}}^{\frac{k}{n}} (k - n\xi) \varphi(\xi) d\xi \\ &= \sum_{k=1}^n (A^n v^n)_k g_k + (A^n v^n)_{k-1} \bar{g}_{k-1} \\ &= -n^2 \sum_{k=1}^n (v_k^n - v_{k-1}^n) ((g_k - g_{k-1}) + (\bar{g}_k - \bar{g}_{k-1})), \end{aligned} \quad (14)$$

where we denote the integrals by g_k and $\bar{g}_{k-1}$, respectively. Note that the boundary term vanish because of the boundary conditions for v^n . The next step is to replace φ by E^n but since the computations differ for v and the piecewise linear $\tilde{v}^n$ we split these terms and obtain the following lemmas.

Lemma 5.1. *For all $n \in \mathbb{N}$ it holds that*

$$V^* \langle A\tilde{v}^n - \iota_n A^n v^n, \tilde{v}^n \rangle_V \leq 0.$$

Proof. Since $\tilde{v}^n$ is piecewise linear we can compute g_k and $\bar{g}_k$, which are

$$g_k = \frac{1}{3n} v_k^n + \frac{1}{6n} v_{k-1}^n \quad \text{and} \quad \bar{g}_k = \frac{1}{6n} v_{k+1}^n + \frac{1}{3n} v_k^n.$$

Obviously, we also have $\tilde{v}^n(\frac{k}{n}) = v_k^n$ and $\tilde{v}^n(\frac{k-1}{n}) = v_{k-1}^n$. Equations (13) and (14) now read as

$$\begin{aligned} V^* \langle A\tilde{v}^n - \iota_n A^n v^n, \tilde{v}^n \rangle_V &= -n \sum_{k=1}^n (v_k^n - v_{k-1}^n)^2 + \frac{2n}{3} \sum_{k=1}^n (v_k^n - v_{k-1}^n)^2 \\ &\quad + \frac{n}{6} \sum_{k=2}^n (v_k^n - v_{k-1}^n)(v_{k-1}^n - v_{k-2}^n) + \frac{n}{6} \sum_{k=1}^{n-1} (v_k^n - v_{k-1}^n)(v_{k+1}^n - v_k^n) \end{aligned}$$

With Cauchy's inequality we can bound the latter terms by the former ones, thus the statement is proven. $\square$

Lemma 5.2. *Let $\varphi \in V$, then*

$$V^* \langle A\tilde{v}^n - \iota_n A^n v^n, \varphi \rangle_V \leq 2 \|\tilde{v}^n\|_V \cdot I_n(\varphi),$$

where

$$I_n(\varphi) := \left(\sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left(\varphi'(\zeta) - \varphi'(\zeta \pm \frac{1}{n}) \right)^2 d\zeta \right)^{\frac{1}{2}}.$$

Proof. The factor with φ in equations (13) and (14) can be written as follows

$$\begin{aligned} & \frac{1}{n} \left(\varphi\left(\frac{k}{n}\right) - \varphi\left(\frac{k-1}{n}\right) \right) - (g_k - g_{k-1}) + (\bar{g}_k - \bar{g}_{k-1}) \\ &= \int_{\frac{k-1}{n}}^{\frac{k}{n}} (n\xi - k + 1) \left[\left(\varphi\left(\frac{k}{n}\right) - \varphi(\xi) \right) - \left(\varphi\left(\frac{k-1}{n}\right) - \varphi\left(\xi - \frac{1}{n}\right) \right) \right] \\ & \quad + (k - n\xi) \left[\left(\varphi(\xi) - \varphi\left(\frac{k-1}{n}\right) \right) - \left(\varphi\left(\xi + \frac{1}{n}\right) - \varphi\left(\frac{k}{n}\right) \right) \right] d\xi =: c_n(\varphi), \end{aligned}$$

because $(n\xi - k + 1) + (k - n\xi) = 1$ and the integrals appearing in g_{k-1} and $\bar{g}_k$ are shifted to the same interval. Rewrite $c_n(\varphi)$ with the fundamental theorem of calculus and use Jensen's inequality to obtain

$$\begin{aligned} c_n(\varphi)^2 &= \left(\int_{\frac{k-1}{n}}^{\frac{k}{n}} (n\xi - k + 1) \int_{\xi}^{\frac{k}{n}} \varphi'(\zeta) - \varphi'\left(\zeta - \frac{1}{n}\right) d\zeta d\xi \right. \\ & \quad \left. + \int_{\frac{k-1}{n}}^{\frac{k}{n}} (k - n\xi) \int_{\frac{k-1}{n}}^{\xi} \varphi'(\zeta) - \varphi'\left(\zeta + \frac{1}{n}\right) d\zeta d\xi \right)^2 \\ &\leq 2n^{-3} \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left(\varphi'(\zeta) - \varphi'\left(\zeta \pm \frac{1}{n}\right) \right)^2 d\zeta. \end{aligned} \tag{15}$$

All there remains to do is an application of the Cauchy-Schwarz inequality. $\square$

Lemma 5.3. *Let $I_n(\varphi)$ be defined as in Lemma 5.2.*

i. If $\varphi \in V$, then $I_n(\varphi) \leq 4\|\varphi\|_V$ and moreover $\lim_{n \rightarrow \infty} I_n(\varphi) = 0$.

ii. If $\varphi \in H^{1+\alpha}(0, 1)$, the explicit rate of convergence is given by $I_n(\varphi) \leq 2^{\alpha+2} n^{-\alpha} \|\varphi\|_{H^{1+\alpha}}$.

Proof. At first, we will consider the second assertion. Thus, let $\varphi \in C^{1+\alpha}([0, 1])$ and denote by φ' its first derivative. Recall the definition of $c_n(\varphi)$ in the proof above. An additional integral in ξ over the same interval yields another n , while we can insert $\varphi'(\xi)$ into the square above. The distance $|\zeta \pm \frac{1}{n} - \xi|$ is of course bounded by $\frac{2}{n}$, hence

$$c_n(\varphi)^2 \leq 2^{2\alpha+3} n^{-3-2\alpha} \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{k-1}{n}}^{\frac{k}{n}} \frac{(\varphi'(\zeta) - \varphi'(\xi))^2}{|\zeta - \xi|^{2\alpha+1}} + \frac{(\varphi'(\xi) - \varphi'(\zeta \pm \frac{1}{n}))^2}{|\zeta \pm \frac{1}{n} - \xi|^{2\alpha+1}} d\zeta d\xi$$

and we have shown *ii.*, since the sum of the latter expression is uniformly bounded by the Sobolev-Slobodeckij semi-norm of φ in $H^{1+\alpha}$.

In a second step, we prove that if $\varphi \in V$, the expression $I_n(\varphi)$ is bounded uniformly in n and we can indeed approximate φ by sufficiently smooth functions in V to obtain the desired convergence. For this purpose, we estimate (15) simply by

$$c_n(\varphi)^2 \leq 8n^{-3} \int_{\frac{k-1}{n}}^{\frac{k}{n}} |\varphi'(\zeta)|^2 + |\varphi'(\zeta \pm \frac{1}{n})|^2 d\zeta,$$

hence $I_n(\varphi) \leq 4\|\varphi\|_V$. Now, let $\varepsilon > 0$ be given. Clearly, $C^{1+\alpha}([0, 1]) \subset V$ densely for $\alpha > 0$, therefore one can find $\varphi_\varepsilon \in C^{1+\alpha}([0, 1])$ such that $\|\varphi - \varphi_\varepsilon\|_V \leq \frac{\varepsilon}{8}$. Furthermore, we can find $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$

$$I_n(\varphi) \leq I_n(\varphi - \varphi_\varepsilon) + I_n(\varphi_\varepsilon) \leq 4\|\varphi - \varphi_\varepsilon\|_V + 2^{\alpha+2}n^{-\alpha}\|\varphi_\varepsilon\|_{H^{1+\alpha}} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon. \quad \square$$

If we turn our focus back on the original error term E_1 and apply the previous lemmas, we have shown that

$$E_1 \leq -2 \int_0^t \|v(s) - \tilde{v}^n(s)\|_V^2 ds + 2 \int_0^t \|\tilde{v}^n(s)\|_V I_n(v(s)) ds. \quad (16)$$

Step 3: Approximation error of the nonlinearities

The nonlinear terms in E_2 are estimated with the (local) Lipschitz conditions (A5) and (A6) together with the uniform a priori estimates on the approximating solutions $\tilde{v}_n$ and $\tilde{w}^n$. The error term in E_2 is

$$\nu^* \langle \Phi(\tilde{v}^n(s), \tilde{w}^n(s)) - \iota_n \Phi^n(v^n(s), w^n(s)), E^n(s) \rangle_\nu$$

which consists of contributing parts of ϕ_1 and ϕ_2 . Since the assumptions on ϕ_2 are much stronger, we exemplarily estimate the one for ϕ_1 in the following.

$$\begin{aligned} & \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left[\phi_1(\xi, \tilde{v}^n(\xi), \tilde{w}^n(\xi)) \right. \\ & \quad \left. - (n\xi - k + 1)\phi_1\left(\frac{k}{n}, v_k^n, w_k^n\right) - (k - n\xi)\phi_1\left(\frac{k-1}{n}, v_{k-1}^n, w_{k-1}^n\right) \right] (v(\xi) - \tilde{v}^n(\xi)) d\xi \\ & \leq \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left[|(n\xi - k + 1)| \cdot |\phi_1(\xi, \tilde{v}^n(\xi), \tilde{w}^n(\xi)) - \phi_1\left(\frac{k}{n}, v_k^n, w_k^n\right)| \right. \\ & \quad \left. + |k - n\xi| \cdot |\phi_1(\xi, \tilde{v}^n(\xi), \tilde{w}^n(\xi)) - \phi_1\left(\frac{k-1}{n}, v_{k-1}^n, w_{k-1}^n\right)| \right] |v(\xi) - \tilde{v}^n(\xi)| d\xi \\ & \leq \sum_{k=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} L \left[(1 + |v_k^n|^{m-1} + |v_{k-1}^n|^{m-1}) \right. \\ & \quad \left. \cdot \left(\frac{1}{n} + |v_k^n - v_{k-1}^n| + |w_k^n - w_{k-1}^n| \right) \right] |v(\xi) - \tilde{v}^n(\xi)| d\xi, \end{aligned}$$

where we used (A3), $|\tilde{v}^n(\xi)| \leq |v_k^n| + |v_{k-1}^n|$ and that $|\tilde{v}^n(\xi) - v_k^n| = (k - n\xi)|v_k^n - v_{k-1}^n|$ as well as $|\tilde{v}^n(\xi) - v_{k-1}^n| = (n\xi - k + 1)|v_k^n - v_{k-1}^n|$. Of course, the same holds for $\tilde{w}^n$. With Cauchy-Schwarz' and Young's inequality we can further estimate

$$\begin{aligned} & \leq \frac{1}{2n^2} L \sum_{k=1}^n (1 + |v_k^n|^{m-1})^2 + \frac{n}{2} L \sum_{k=1}^n \left(\frac{1}{n} + |v_k^n - v_{k-1}^n| + |w_k^n - w_{k-1}^n| \right)^2 \int_{\frac{k-1}{n}}^{\frac{k}{n}} |v(\xi) - \tilde{v}^n(\xi)|^2 d\xi \\ & \leq \frac{1}{n} L \left(1 + l_{m+1}^n(v^n) \right) + \frac{3}{2} L (1 + \|\tilde{v}^n\|_V^2 + \|\tilde{w}^n\|_V^2) \|v - \tilde{v}^n\|_H^2, \end{aligned}$$

since $2(m-1) \leq m+1$ for $m \leq 3$ and the gradient of the linear interpolation is given by these finite differences, see Proposition 4.3. For simplicity we estimate ϕ_2 's part of the error in the

same way, hence

$$E_2 \leq 6L \int_0^t (1 + \|\tilde{v}^n(s)\|_V^2 + \|\tilde{w}^n(s)\|_V^2) \|E^n(s)\|_{\mathcal{H}}^2 ds + \frac{4}{n} L \int_0^t (1 + l_{m+1}^n(v^n(s))) ds. \quad (17)$$

Step 4: Approximation of the covariance operator

The Itô correction term gives us a Hilbert-Schmidt norm which, in our case, is given by the L^2 -norm of the associated kernel. Thus, we can write this part of the error term E_4 similar to the one of the nonlinear part in Step 3. Recall that $P_n h = (\langle h, \sqrt{n} 1_{[\frac{l-1}{n}, \frac{l}{n})} \rangle)_{1 \leq l \leq n}$.

$$\begin{aligned} & \|B(\tilde{v}^n) \sqrt{Q} - \iota_n b^n(v^n) \sqrt{Q^n} P_n\|_{\text{HS}}^2 \\ &= \int_0^1 \int_0^1 \left(b(\xi, \tilde{v}^n(\xi)) q(\xi, \zeta) - \sum_{k=1}^n 1_{[\frac{k-1}{n}, \frac{k}{n})}(\xi) \right. \\ & \quad \cdot \left. \left[(n\xi - k + 1) b(\frac{k}{n}, v_k^n) \sum_{l=1}^n q_{k,l}^n 1_{[\frac{l-1}{n}, \frac{l}{n})}(\zeta) + (k - n\xi) b(\frac{k-1}{n}, v_{k-1}^n) \sum_{l=1}^n q_{k-1,l}^n 1_{[\frac{l-1}{n}, \frac{l}{n})}(\zeta) \right] \right)^2 d\zeta d\xi. \end{aligned}$$

A closer look reveals that the error consists of parts S_1 contributed by the approximation of b and S_2 by the approximation of the kernel q . Therefore we split the norm square into two parts where the first one is

$$2 \sum_{k=1}^n \int_0^1 \int_{\frac{k-1}{n}}^{\frac{k}{n}} \left(b(\xi, \tilde{v}^n(\xi)) - (n\xi - k + 1) b(\frac{k}{n}, v_k^n) - (k - n\xi) b(\frac{k-1}{n}, v_{k-1}^n) \right)^2 q(\xi, \zeta)^2 d\zeta d\xi.$$

Similar to Step 3 for the nonlinear drift term, since b is Lipschitz continuous,

$$\left(b(\xi, \tilde{v}^n(\xi)) - (n\xi - k + 1) b(\frac{k}{n}, v_k^n) - (k - n\xi) b(\frac{k-1}{n}, v_{k-1}^n) \right)^2 \leq \frac{1}{2} \left(\frac{1}{n^2} + |v_k^n - v_{k-1}^n|^2 \right)$$

and the summation over n yields

$$S_1 \leq \frac{1}{n} \text{tr } Q (1 + \|\tilde{v}^n\|_V^2).$$

The second error S_2 term due to the approximation of the covariance kernel is given by

$$4 \sum_{k,l=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} \left(b(\frac{k}{n}, v_k^n) (q(\xi, \zeta) - q_{k,l}^n) \right)^2 + \left(b(\frac{k-1}{n}, v_{k-1}^n) (q(\xi, \zeta) - q_{k-1,l}^n) \right)^2 d\zeta d\xi,$$

where we already used $(n\xi - k + 1) \leq 1$ as well as $(k - n\xi) \leq 1$. Exemplarily we do the estimate for the part with $q_{k,l}^n$ but note that the procedure remains the same in the one with $q_{k-1,l}^n$ except for a constant. Since b is bounded, we obtain

$$\begin{aligned} & \leq 4 \sum_{k,l=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} (q(\xi, \zeta) - q_{k,l}^n)^2 d\zeta d\xi \\ &= 4 \sum_{k,l=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} \left(q(\xi, \zeta) - n^2 \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} q(\rho, \eta) d\rho d\eta \right)^2 d\zeta d\xi \\ & \leq 4n^2 \sum_{k,l=1}^n \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} \int_{\frac{k-1}{n}}^{\frac{k}{n}} \int_{\frac{l-1}{n}}^{\frac{l}{n}} |q(\xi, \zeta) - q(\rho, \eta)|^2 d\xi d\zeta d\rho d\eta. \end{aligned}$$

Since $|\xi - \rho| \leq \frac{1}{n}$ as well as $|\zeta - \eta| \leq \frac{1}{n}$ we can expand this expression to the Sobolev-Slobodeckij semi-norm of q .

$$\begin{aligned} &\leq 2^{\theta+3} n^{-2\theta} \iiint_0^1 \frac{|q(\xi, \zeta) - q(\rho, \eta)|^2}{(|\xi - \rho|^2 + |\zeta - \eta|^2)^{\theta+1}} d\xi d\zeta d\rho d\eta \\ &\leq 2^{\theta+3} n^{-2\theta} \|q\|_{H^\theta}^2. \end{aligned}$$

In summary, we have shown that (the modified constants are due to the omitted terms)

$$E_4 \leq 2c_\theta \operatorname{tr}(-A)^\theta Q \int_0^t \|E^n(s)\|_{\mathcal{H}}^2 ds + \frac{2}{n} \operatorname{tr} Q \int_0^t (1 + \|\tilde{v}^n(s)\|_V^2) ds + 2^{3\theta+5} n^{-2\theta} \|q\|_{H^\theta}^2 \cdot t \quad (18)$$

Step 5: Control over the supremum

The four previous steps allow a first estimate non-uniform in t . For this purpose define the processes

$$L^n(t) := 6L(1 + \|\tilde{v}^n(t)\|_V^2 + \|\tilde{w}^n(t)\|_V^2) + 2c_\theta \operatorname{tr}(-A)^\theta Q$$

and

$$\begin{aligned} K^n(t) := \frac{1}{n} \Big(4L(1 + l_{m+1}^n(v^n(t))) + 2 \operatorname{tr} Q(1 + \|\tilde{v}^n(t)\|_V^2) \\ + 2^{3\theta+5} \|q\|_{H^\theta}^2 \Big) + 2\|\tilde{v}^n(t)\|_V I_n(v(t)). \end{aligned}$$

Now we apply Itô's formula with the function $f(x) = x^p$, $p \geq 1$ for the square of the $\mathcal{H}$ -norm of the error $E^n(t)$. By Steps 1-4

$$\begin{aligned} \|E^n(t)\|_{\mathcal{H}}^{2p} &\leq \|E^n(0)\|_{\mathcal{H}}^{2p} + p \int_0^t L^n(s) \|E^n(s)\|_{\mathcal{H}}^{2p} ds + p \int_0^t K^n(s) \|E^n(s)\|_{\mathcal{H}}^{2p-2} ds \\ &\quad + p \int_0^t \|E^n(s)\|_{\mathcal{H}}^{2p-2} dM(s) + \frac{p(p-1)}{2} \int_0^t \|E^n(s)\|_{\mathcal{H}}^{2p-4} d\langle M \rangle_s, \end{aligned}$$

where the local martingale $M(t)$ is given by

$$M(t) := 2 \int_0^t \langle v(s) - \tilde{v}^n(s), (B(v(s))\sqrt{Q} - \iota_n b^n(v^n(s))\sqrt{Q^n} P_n) dW(s) \rangle.$$

Its quadratic variation is simply

$$\langle M \rangle_t = \int_0^t \|v(s) - \tilde{v}^n(s)\|_H^2 \|B(v(s))\sqrt{Q} - \iota_n b^n(v^n(s))\sqrt{Q^n} P_n\|_{\text{HS}}^2 ds.$$

Obviously, there appears exactly the same term as in E_4 which we already estimated point-wise in t . Thus, use (18) and Itô's product rule to conclude

$$\begin{aligned} \sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} &\leq \|E^n(0)\|_{\mathcal{H}}^{2p} + \int_0^T e^{-\int_0^s L_p^n(r) dr} K_p^n(t) \|E^n(t)\|_{\mathcal{H}}^{2p-2} dt \\ &\quad + \sup_{t \in [0, T]} \int_0^t e^{-\int_0^s L_p^n(r) dr} \|E^n(s)\|_{\mathcal{H}}^{2p-2} dM(s). \end{aligned} \quad (19)$$

The processes $L_p^n(t)$ and $K_p^n(t)$ are given by

$$L_p^n(t) := pL^n(t) + p(p-1)c_\theta \operatorname{tr}(-A)^\theta Q$$

and

$$K_p^n(t) := pK^n(t) + \frac{1}{n}p(p-1)\left(\operatorname{tr} Q(1 + \|\tilde{v}^n(t)\|_V^2) + 2^{3\theta+4}\|q\|_{H^\theta}^2\right).$$

The first integral in (19) can be estimated with Young's inequality as follows

$$\begin{aligned} \int_0^T e^{-\int_0^t L_p^n(s) ds} K_p^n(t) \|E^n(t)\|_{\mathcal{H}}^{2p-2} dt &\leq \sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p-2} \int_0^T K_p^n(t) dt \\ &\leq \frac{p-1}{p} \sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} + \frac{1}{p} \left(\int_0^T K_p^n(t) dt \right)^p \end{aligned}$$

Now, take the expectation in (19) and the Burkholder-Davis-Gundy inequality bounds the supremum of the stochastic integral from above by its quadratic variation, more precisely

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} \right] &\leq p \mathbb{E} \left[\|E^n(0)\|_{\mathcal{H}}^{2p} \right] + \mathbb{E} \left[\left(\int_0^T K_p^n(t) dt \right)^p \right] \\ &\quad + 3p \mathbb{E} \left[\left(\int_0^T e^{-2 \int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{4p-4} d\langle M \rangle_t \right)^{\frac{1}{2}} \right]. \end{aligned} \quad (20)$$

The quadratic variation is again estimated as above in terms of the left hand side and $K_p^n(t)$. This is done with Young's inequality as follows

$$\begin{aligned} 3p \mathbb{E} \left[\left(\int_0^T e^{-2 \int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{4p-4} d\langle M \rangle_t \right)^{\frac{1}{2}} \right] &\leq \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\frac{2p-1}{2p} \int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p-1} \right. \\ &\quad \cdot \left. \left(\int_0^T e^{-\frac{1}{p} \int_0^t L_p^n(s) ds} \|B(v(t))\sqrt{Q} - \iota_n b^n(v^n(t))\sqrt{Q^n} P_n\|_{\text{HS}}^2 dt \right)^{\frac{1}{2}} \right] \\ &\leq 3(6p-3)^{2p-1} \mathbb{E} \left[\left(\int_0^T e^{-\frac{1}{p} \int_0^t L_p^n(s) ds} \|B(v(t))\sqrt{Q} - \iota_n b^n(v^n(t))\sqrt{Q^n} P_n\|_{\text{HS}}^2 dt \right)^p \right] \\ &\quad + \frac{1}{2} \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} \right] \end{aligned}$$

The latter is absorbed by the left hand side and the former is again essentially E_4 which we already estimated in Step 4. With (18) and (20) follows

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} \right] &\leq 2p \mathbb{E} \left[\|E^n(0)\|_{\mathcal{H}}^{2p} \right] + C_3 \mathbb{E} \left[\left(\int_0^T K_p^n(t) dt \right)^p \right] \\ &\quad + C_4 \mathbb{E} \left[\int_0^T \sup_{s \in [0, t]} e^{-\int_0^s L_p^n(r) dr} \|E^n(s)\|_{\mathcal{H}}^{2p} dt \right], \end{aligned}$$

where $C_3 := 6(6p - 3)^{2p-1}2^p + 2$ and $C_4 := 12(6p - 3)^{2p-1}2^p c_\theta \operatorname{tr}(-A)^\theta Q$. Now, Gronwall's inequality implies for all $p \geq 1$

$$\mathbb{E} \left[\sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} \right] \leq e^{C_4 T} \left(2p \mathbb{E} [\|E^n(0)\|_{\mathcal{H}}^{2p}] + C_3 \mathbb{E} \left[\left(\int_0^T K^n(t) dt \right)^p \right] \right). \quad (21)$$

This preliminary error estimate yields the desired one via Hölder's inequality, since we have (21) for every $p \geq 1$ and can control the exponential by Proposition 4.3. Therefore, we have for $p \geq 1$

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} \|E^n(t)\|_{\mathcal{H}}^p \right]^{\frac{1}{p}} &\leq \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\frac{1}{2} \int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^p e^{\frac{1}{2} \int_0^t L_p^n(s) ds} \right]^{\frac{1}{p}} \\ &\leq \mathbb{E} \left[\sup_{t \in [0, T]} e^{-\int_0^t L_p^n(s) ds} \|E^n(t)\|_{\mathcal{H}}^{2p} \right]^{\frac{1}{2p}} \mathbb{E} \left[e^{\int_0^T L_p^n(t) dt} \right]^{\frac{1}{2p}} \\ &\leq C_{4.3}^{\frac{1}{2p}} e^{C_4 T} \left(2p \mathbb{E} [\|E^n(0)\|_{\mathcal{H}}^{2p}] + C_3 \mathbb{E} \left[\left(\int_0^T K_p^n(t) dt \right)^p \right] \right)^{\frac{1}{2p}}. \quad (22) \end{aligned}$$

It remains to study the convergence of K^n to 0. For this purpose, we fix $p = 1$ at first. Then $K_1^n(t) = K^n(t)$ and by Proposition 4.3

$$\mathbb{E} \left[\int_0^T K^n(t) dt \right] \leq Cn^{-1} + 2\mathbb{E} \left[\int_0^T \|\tilde{v}^n(t)\|_V I_n(v(t)) dt \right]$$

Since by Lemma 5.2 $I_n(v(t)) \leq 4\|v(t)\|_V$ and both v and $\tilde{v}^n$ are (uniformly) bounded in $\mathcal{L}^2(\Omega, \mathcal{F}, \mathbb{P}; L^2([0, T], V))$ we can use Lebesgue's dominated convergence theorem to deduce

$$\mathbb{E} \left[\int_0^T K^n(t) dt \right] \rightarrow 0, \quad \text{hence} \quad \mathbb{E} \left[\sup_{t \in [0, T]} \|E^n(t)\|_{\mathcal{H}} \right] \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

by Lemma 5.2. Thus, the exponential moment estimates carry over to the limit v and we can apply the dominated convergence theorem in cases $p > 1$ which concludes the first assertion of Theorem 3.1. If moreover $v \in \mathcal{L}^{p^*}(\Omega, \mathcal{F}, \mathbb{P}; L^2([0, T], H^{1+\alpha}))$ for some $\alpha > 0$ and $p^* > 1$, then again by Lemma 5.2

$$\begin{aligned} \mathbb{E} \left[\left(\int_0^T K^n(t) dt \right)^p \right] &\leq Cn^{-p} + C' \mathbb{E} \left[\left(\int_0^T \|\tilde{v}^n(t)\|_V^2 dt \right)^{\frac{p}{2}} \left(\int_0^T I_n(v(t))^2 dt \right)^{\frac{p}{2}} \right] \\ &\leq Cn^{-p} + C' \mathbb{E} \left[\left(\int_0^T \|\tilde{v}^n(t)\|_V^2 dt \right)^{\frac{pp^*}{2(p^*-p)}} \right]^{\frac{p^*-p}{p^*}} \mathbb{E} \left[\left(\int_0^T I_n(v(t))^2 dt \right)^{\frac{p^*}{2}} \right]^{\frac{p}{p^*}} \\ &\leq Cn^{-p} + C'' n^{-\alpha p} \mathbb{E} \left[\left(\int_0^T \|v(t)\|_{H^{1+\alpha}}^2 dt \right)^{\frac{p^*}{2}} \right]^{\frac{p}{p^*}}, \end{aligned}$$

hence the second assertion is proven for all $p < p^*$. $\square$

6 Applications

As an application for our results in Theorem 3.1, we consider the spatially extended FitzHugh-Nagumo system with noise. Originally, this was stated by FitzHugh [3] as a system of ODEs simplifying the famous Hodgkin-Huxley model [8] for the generation of an action potential in a neuron in terms of a voltage variable v and a so-called recovery variable w . Its spatially extended version is a model for the propagation of the action potential in the axon of a neuron. See e.g. the monographs [2, 12] for more details on the deterministic case. Now consider this system subject to external noise only in the voltage variable v . Together with the original parameters from [4] this reads as

$$\begin{aligned} \dot{v}(t, \xi) &= \partial_{\xi}^2 v(t, \xi) + v(t, \xi) - \frac{1}{3}v(t, \xi)^3 - w(t, \xi) + \eta(t, \xi), \\ \dot{w}(t, \xi) &= 0.08(v(t, \xi) - 0.8w(t, \xi) + 0.7), \quad t \geq 0, \xi \in (0, L) \end{aligned} \quad (23)$$

equipped with homogeneous Neumann boundary conditions in 0 and L . One can immediately see, that (23) is of the type (1). The first mathematical rigorous analysis of this equation in the context of mild solutions can be found in [1]. It has been observed in e.g. [18] that this system has traveling pulse solutions (Figure 1) which may break down due to the influence of the external noise, hence there is no transmission of the signal from 0 to L . This phenomenon is usually referred to as the propagation failure and one is interested in calculating its probability depending on the strength of the external noise, see [18] for a heuristic approach.




Figure 1: A traveling pulse solution of (23) obtained by the finite difference scheme propagating along $(0, L)$ at three different times (solid, dashed, dotted).

An appropriate estimator detecting the event of a propagation failure is given by the integral

$$\Phi(v) := \int_0^L v(\xi) - v_* \, d\xi,$$

which significantly differs in cases with or without the traveling pulse based on the observation from Figure 1. Here, $v_* \approx -1.1994$ is the voltage component of the unique real equilibrium point for the drift in (23). Given Φ , the event of a propagation failure can be defined by

$$\Phi(v(t)) \leq \kappa \quad \text{for some } T_0 \leq t \leq T,$$

for some appropriate threshold $\kappa > 0$ and some initializing time T_0 . The quantity of interest is the probability

$$\mathbb{P}_Q \left[\min_{T_0 \leq t \leq T} \Phi(v(t)) \leq \kappa \right] =: p_{Q,\kappa}$$

of propagation failure depending on the noise covariance Q . Essentially we have to estimate the parameter of a Bernoulli distributed random variable. The sample average

$$\hat{p}_{Q,\kappa}(\hat{v}) := \frac{1}{m} \sum_{k=1}^m 1_{\{\min_{T_0 \leq t \leq T} \Phi(v^{(k)}(t)) \leq \kappa\}}$$

based on m iid copies $\hat{v} := (v^{(k)})_{1 \leq k \leq m}$ is a natural estimator. Since the solution v is not given explicitly, we can approximate $\hat{p}_{Q,\kappa}$ based on independent numerical observations $\hat{v}^n := (\tilde{v}^{n,(k)})_{1 \leq k \leq m}$, as independent realizations of $\tilde{v}^n$ only. Besides the statistical Monte-Carlo error, there appears additional uncertainty due to the approximation of the exact solution. Theorem 3.1 now allows to quantify this.

Corollary 6.1. *Let $\hat{p}_{Q,\kappa}^n := \hat{p}_{Q,\kappa}(\hat{v}^n)$ and $\varepsilon > 0$ be a priori given. Furthermore, assume that the solution $v \in \mathcal{L}^p(\Omega, \mathcal{F}, \mathbb{P}; L^2([0, T], H^2(0, 1)))$ for some $p > 2$. Then, a confidence interval for the estimation $\hat{p}_{Q,\kappa}^n$ of $p_{Q,\kappa}$ with confidence level α is given by $[\hat{p}_{Q,\kappa}^n - \gamma, \hat{p}_{Q,\kappa}^n + \gamma]$ where*

$$\gamma = (\alpha m)^{-\frac{1}{2}} \left(1 + 4\varepsilon^{-2} C n^{-1} \right)^{\frac{1}{2}}.$$

Remark. The additional regularity of the solution may be obtained in the context of mild solutions as in [1] if Q is sufficiently regular, since the heat semigroup in the equation for v is analytic and therefore maps H to $D(A) = H^2(0, 1)$. However, this needs further investigation.

Proof. We can estimate the probability of $p_{Q,\kappa}$ being outside of an interval of size γ around the estimator $\hat{p}_{Q,\kappa}^n$ by Chebychev's inequality with

$$\begin{aligned} \mathbb{P}_Q [|p_{Q,\kappa} - \hat{p}_{Q,\kappa}^n| > \gamma] &= \mathbb{P}_Q [|p_{Q,\kappa} - \hat{p}_{Q,\kappa} + \hat{p}_{Q,\kappa} - \hat{p}_{Q,\kappa}^n| > \gamma] \\ &\leq \mathbb{P}_Q [|p_{Q,\kappa} - \hat{p}_{Q,\kappa}| > \frac{\gamma}{2}] + \mathbb{P}_Q [|\hat{p}_{Q,\kappa} - \hat{p}_{Q,\kappa}^n| > \frac{\gamma}{2}] \\ &\leq \frac{1}{\gamma^2 m} + \mathbb{P}_Q \left[\sum_{k=1}^m 1_{\{\sup_{t \in [0, T]} |\Phi(v^{(k)}(t)) - \Phi(\tilde{v}^{n,(k)}(t))| > \varepsilon\}} > m \frac{\gamma}{2} \right] \\ &\leq \frac{1}{\gamma^2 m} + \frac{4}{\gamma^2 \varepsilon^2 m} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\Phi(v^{(k)}(t)) - \Phi(\tilde{v}^{n,(k)}(t)) \right)^2 \right]. \end{aligned}$$

The additional uncertainty due to the approximation of the exact solution is the mean squared error of $\Phi(\tilde{v}^n(t))$.

$$\begin{aligned} \mathbb{E} \left[\sup_{t \in [0, T]} \left(\Phi(v(t)) - \Phi(\tilde{v}^n(t)) \right)^2 \right] &= \mathbb{E} \left[\sup_{t \in [0, T]} \left(\int_0^L v(t, \xi) - v_* d\xi - \int_0^L \tilde{v}^n(t, \xi) - v_* d\xi \right)^2 \right] \\ &\leq L \mathbb{E} \left[\sup_{t \in [0, T]} \|v(t) - \tilde{v}^n(t)\|_H^2 \right] \leq L \mathbb{E} \left[\sup_{t \in [0, T]} \|E^n(t)\|_{\mathcal{H}}^2 \right]. \end{aligned}$$

Theorem 3.1 now implies the convergence rate of n^{-1} and we set

$$\alpha = \frac{1}{\gamma^2 m} + \frac{4L}{\gamma^2 \varepsilon^2 m} C n^{-1}. \quad \square$$

References

- [1] S. Bonaccorsi and E. Mastrogiacomo. Analysis of the Stochastic FitzHugh-Nagumo System. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 11(03):427–446, 2008.
- [2] G. B. Ermentrout and D. H. Terman. *Mathematical Foundations of Neuroscience*, volume 35. Springer, 2010.
- [3] R. FitzHugh. Impulses and Physiological States in Theoretical Models of Nerve Membrane. *Biophys. J.*, 1:445–466, 1961.
- [4] R. FitzHugh. Mathematical Models of Excitation and Propagation in Nerve. In *Biological Engineering*. McGrawHill, New York, 1969.
- [5] I. Gyöngy. Lattice Approximations for Stochastic Quasi-Linear Parabolic Partial Differential Equations Driven by Space-Time White Noise II. *Potential Anal.*, 11(1):1–37, 1999.
- [6] I. Gyöngy and A. Millet. On Discretization Schemes for Stochastic Evolution Equations. *Potential Anal.*, 23(2):99–134, 2005.
- [7] I. Gyöngy and A. Millet. Rate of Convergence of Space Time Approximations for Stochastic Evolution Equations. *Potential Anal.*, 30(1):29–64, 2009.
- [8] A. L. Hodgkin and A. F. Huxley. A Quantitative Description of Membrane Current and its Application to Conduction and Excitation in Nerve. *J. Physiol.*, 117:500–544, 1952.
- [9] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong and Weak Divergence in Finite Time of Euler’s Method for Stochastic Differential Equations With Non-Globally Lipschitz Continuous Coefficients. *Proc. R. Soc. A*, 467(2130):1563–1576, 2011.
- [10] M. Hutzenthaler, A. Jentzen, and P. E. Kloeden. Strong Convergence of an Explicit Numerical Method for SDEs With Nonglobally Lipschitz Continuous Coefficients. *Ann. Appl. Probab.*, 22(4):1611–1641, 2012.
- [11] A. Jentzen. Pathwise Numerical Approximation of SPDEs with Additive Noise under Non-Global Lipschitz Coefficients. *Potential Anal.*, 31(4):375–404, 2009.
- [12] J. Keener and J. Sneyd. *Mathematical Physiology: I: Cellular Physiology*, volume 1. Springer, 2008.
- [13] D. Liu. Convergence of the Spectral Method for Stochastic Ginzburg-Landau Equation Driven by Space-Time White Noise. *Commun. Math. Sci.*, 1(2):361–375, 2003.
- [14] W. Liu and M. Röckner. SPDE in Hilbert Space with Locally Monotone Coefficients. *J. Funct. Anal.*, 259(11):2902–2922, 2010.

- [15] R. Pettersson and M. Signahl. Numerical Approximation for a White Noise Driven SPDE with Locally Bounded Drift. *Potential Anal.*, 22(4):375–393, 2005.
- [16] C. Prévôt and M. Röckner. *A Concise Course on Stochastic Partial Differential Equations*, volume 1905 of *Lecture Notes in Mathematics*. Springer, Berlin, 2007.
- [17] H. Triebel. *Interpolation Theory, Function Spaces, Differential Operators*, volume 18 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1978.
- [18] H. C. Tuckwell. Analytical and Simulation Results for the Stochastic Spatial FitzHugh-Nagumo Model Neuron. *Neural Comput.*, 20(12):3003–3033, 2008.